



Assessment of Methods to Quantify Livestock Associated MRSA in Pig Herds

Hansen, Julie Elvekjær; Sørensen, Anna Irene Vedel; Espinosa-Gongora, Carmen; Larsen, A. R.; Larsen, J.; Skov, R.; Pedersen, Karl

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hansen, J. E., Sørensen, A. I. V., Espinosa-Gongora, C., Larsen, A. R., Larsen, J., Skov, R., & Pedersen, K. (2015). *Assessment of Methods to Quantify Livestock Associated MRSA in Pig Herds*. Poster session presented at 4th ASM-ESCMID Conference on Methicillin-resistant Staphylococci in Animals: Veterinary and Public Health Implications, Chicago, Illinois, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Methods to quantify livestock associated MRSA in pig herds

Julie Elvekjær Hansen¹, Anna Irene Vedel Sørensen¹, Carmen Espinosa-Gongora², Anders Rhod Larsen³, Jesper Larsen³, Robert Skov³ and Karl Pedersen¹

¹Technical University of Denmark, National Veterinary Institute, Frederiksberg, Denmark, ²University of Copenhagen, Department of Veterinary Disease Biology, Frederiksberg, Denmark, ³Statens Serum Institut, Copenhagen, Denmark.

Background A high prevalence of LA-MRSA, especially in the pig production, has been reported from many countries. However, there is a lack of knowledge about the actual LA-MRSA loads on infected farms. The overall MRSA level is low in Denmark and if the reservoir of LA-MRSA starts to spread into the community, the MRSA level in general could increase. We do not believe it is possible to eliminate LA-MRSA in the pig production, but we have an ambition of being able to lower the level. In order to try and control LA-MRSA, it is highly relevant to be able to quantify LA-MRSA in the farm environment and on livestock which would enable us to rank risk factors in the production chain and to assess the effect of possible intervention strategies.

Purpose The present study was undertaken to assess methods to quantify LA-MRSA on animals and in the environment in pig herds. In addition, the role of flies as potential environmental carriers of LA-MRSA was investigated.

Method

Air samples on MRSA 2


Direct plating of swabs on MRSA 2

Colony count


Cfu determination

Results

Method verification



Airborne LA-MRSA



Load in weaning pigs (n=15)





Table 1. Proportions of variance

Source	Nasal swabs		Skin swabs	
	Variance	%	Variance	%
Bio. replicate	0.37	24	0.08	31
Pig	1.11	72	0.13	53
Residual	0.06	4	0.04	15
Total	1.54	100	0.25	100

CFU based on counts from 10⁻⁹, where the 10⁻¹ counts were zero

Nasal swabs seemed to cause greater variation in the results than use of skin swabs and a larger proportion of the variance could be explained by the difference between pigs.



The Sartorius air sampler yielded the most stable detection level of LA-MRSA and the increasing raw counts detected corresponded better with increasing air volumes.

Table 2. Range of MRSA loads.

	Range of MRSA load
Nose	6.6- $\times 10^1$ -3.9 $\times 10^4$ cfu/swab
Skin	1.1 $\times 10^1$ -2.6 $\times 10^5$ cfu/swab
MD8	4.1 $\times 10^2$ -6.3 $\times 10^2$ cfu/m ³

Loads in the 15 weaning pigs supported the results seen in the method verification. Air load and skin load seemed similar while nasal load was higher. MRSA was found on flies, in pools of 5 and individually, after enrichment.

Conclusion Fast quantification of the animal MRSA load in pig herds is possible by direct plating of either nasal or skin swab samples. The Sartorius MD8 air sampler provides fast environmental quantification and it was possible to detect MRSA in flies with animal contact demonstrating their potential as environmental MRSA carriers.

